

Buckeye® 16% Complete Calf Ration

40350

TYPE OF FEED Complete

FORM OF FEED: Texturized

PACKAGING: 50 lb Paper Bag

SPECIES: Calves

AGES: Up to 300 lb



FEATURES:

- Complete pellets that contain the required fiber so that little or no hay is recommended
- Excellent calf starter where calves are kept in housing facilities with manure handling systems which cannot handle hay
- Excellent for outside calf hutches where labor is limited and hay is difficult to feed with considerable waste
- This feed will not set-up in the winter nor attract flies in the summer

GUARANTEED ANALYSIS

Crude Protein, min.	16.0%
Crude Fat, min.	3.00%
Crude Fiber, max.	7.00%
Acid detergent fiber, max.	9.00%
Calcium, min.	0.70%
Calcium, max.	1.20%
Phosphorus, min.	0.50%
Selenium, min.	0.50 ppm
Vitamin A, min.	5000 IU/lb
Vitamin D, min.	1250 IU/lb
Vitamin E, min.	15 IU/lb

INGREDIENTS: Steam Crimped Corn, Oats, Cracked Corn, Soybean Meal, Wheat Middlings, Heat Processed Soybeans Corn Distillers Dried Grains, Soybean Hulls, Cane Molasses, Vegetable Oil, Calcium Carbonate, Calcium Phosphate, Salt, Potassium Chloride, Potassium Sulfate, Magnesium Oxide, Magnesium Sulfate, Manganese Oxide, Cobalt Carbonate, Sodium Selenite, Vitamin A Supplement, Vitamin D-3 Supplement, Vitamin E Supplement, Riboflavin, d-Calcium Pantothenate, Niacin, Vitamin B-12 Supplement, Choline Chloride, Dried Saccharomyces cerevisiae Fermentation Solubles, Brewers Dried Yeast.

FEEDING DIRECTIONS:

	LBS.	LBS
CALF WEIGHT	CALF STARTER	EXCELLENT FORAGE
100-200	1.7 to 3.4	1.0 to 2.5
200-300	3.4 to 5.1	3.0 to 4.5

FEEDING DIRECTIONS: Start calves on Buckeye 16% Complete Calf Ration as early as possible. Feed Buckeye 16% Complete Calf Ration at the rate of 1.7 lb. per 100 lb. of body weight. This product contains yeast products, which will enhance intake as well as partially control pathogenic microorganism. This was added as a replacement for medication based on University studies.

Offer clean, fresh water at all times. Excellent quality hay should be fed from the second week of life.

CAUTION: Never feed more than 60% of the total ration as grain.